

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number  
**WO 03/087032 A1**

- (51) International Patent Classification<sup>7</sup>: C07C 69/734, 259/10, A01N 37/50, C07C 67/343
- (21) International Application Number: PCT/EP03/03784
- (22) International Filing Date: 11 April 2003 (11.04.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
MI2002A000814 17 April 2002 (17.04.2002) IT
- (71) Applicant (for all designated States except US): **ISAGRO RICERCA S.R.L.** [IT/IT]; Via Felice Casati, 20, I-20124 Milan (IT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **VENTURINI, Isabella** [IT/IT]; Via Cavour, 29, I-28062 Cameri-Novara (IT). **BETTARINI, Franco** [IT/IT]; Via Cadore, 4/B, I-28100 Novara (IT). **CASTORO, Paolo** [IT/IT]; Via Rolle 6, I-13100 Vercelli (IT). **CIAPESSONI, Alessandro** [IT/IT]; Via Bologna, 4, I-27100 Pavia (IT). **GUSMEROLI, Marilepa** [IT/IT]; Via Don, Valentini, 20, I-20052 Monza-Milan (IT). **MEAZZA, Giovanni** [IT/IT]; Via G. Pastore, 30, I-21047 Saronno-Varese (IT). **PORTOSO, Domenico** [IT/IT]; Via Marchesi, 9, I-26900 Lodi (IT). **SARGIOTTO, Chiara** [IT/IT]; Corso Francia, 212, I-10146 Torino (IT).
- (74) Agents: **DE GREGORI, Antonella et al.**; Ing. Barzano' & Zanardo Milano S.p.A., Via Borgonuovo, 10, I-20121 Milan (IT).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW); Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM); European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR); OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
  - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANALOGOUS COMPOUNDS OF STROBILURINES AND THEIR USE AS ACARICIDES AND INSECTICIDES



(I)

(57) Abstract: Compounds are described, having general formula (I) and their use as acaricides, insecticides and/or fungicides.

**THIS PAGE BLANK (USPTO)**

10/510383

DT04 Rec'd PCT/PTO 05 OCT 2004

ANALOGOUS COMPOUNDS OF STROBILURINES AND THEIR USE AS ACARICIDES AND  
INSECTICIDES

5

The present invention relates to new analogous com-  
10 pounds of strobilurines. More specifically, the present  
invention relates to new analogous compounds of strobi-  
lurines having a high acaricidal and insecticidal activ-  
ity and their use as acaricides and insecticides.

Analogous products of strobilurines with a fungi-  
15 cidal activity are described in patent applications EP  
178826, EP 226917, EP 253213, EP 278595, EP 398692.


Analogous products of strobilurines with an acarici-  
dal and insecticidal activity are described in patent ap-  
plications EP 242081, EP 299694, EP 335519. In particu-  
20 lar, this latter document describes benzyl-phenyl ethers  
with an acaricidal and insecticidal activity character-  
ized by the presence of the methoxyacrylate group in po-  
sition 2 of the benzyl group and not more than two sub-  
stituents on the phenol ring.

25 The acaricidal/insecticidal activity of the com-

pounds described in the above documents, however, are moderate and cannot be used for practical purposes.

The applicant has now found that in the series of benzyl-phenyl ethers, whose molecular structure is represented by general formula (I), a surprisingly improved acaricidal and insecticidal activity is obtained when a substituent R is present on the phenol ring, as specified below, and at least two of the other four positions are occupied by halogen atoms.

10        An object of the present invention therefore relates to compounds having general formula (I)



(I)

wherein:

- 15    - a group selected from X<sub>1</sub>, X<sub>2</sub> and X<sub>3</sub> represents an R group;
- X<sub>4</sub> and X<sub>5</sub> and two of the remaining X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> groups represent a hydrogen atom or a halogen atom, on the condition that at least two of said groups represent a halogen atom;
- 20

- R represents a C<sub>1</sub>-C<sub>12</sub> alkyl or haloalkyl group; a C<sub>1</sub>-C<sub>12</sub> alkoxy or alkylthio group optionally substituted by: halogen atoms, cyano groups, C<sub>1</sub>-C<sub>6</sub> alkoxy groups optionally halogenated, C<sub>2</sub>-C<sub>10</sub> alkoxyalkoxy groups optionally halogenated, NH<sub>2</sub> groups optionally substituted by C<sub>1</sub>-C<sub>6</sub> alkyl groups optionally halogenated, C<sub>3</sub>-C<sub>12</sub> trialkyl silyl groups, aryloxy or heteroaryloxy groups, in turn optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl groups optionally halogenated, C<sub>1</sub>-C<sub>6</sub> alkoxy groups optionally halogenated, nitro groups, cyano groups; a C<sub>2</sub>-C<sub>12</sub> alkenyloxy or alkenylthio group optionally substituted by halogen atoms, cyano groups, aryl groups or heteroaryl groups, in turn optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, halo-alkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a C<sub>3</sub>-C<sub>12</sub> alkynyloxy or alkynylthio group optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkoxy or haloalkoxy groups, aryl or heteroaryl groups, in turn optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, haloalkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a linear or branched C<sub>3</sub>-C<sub>12</sub> alkoxyiminoalkylidenoxy or alkoxyiminoalkylidenthio group; a C<sub>3</sub>-C<sub>8</sub> cycloalkoxy or cycloalkylthio group optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, haloalkyl, alkoxy, haloalkoxy groups; a C<sub>4</sub>-C<sub>12</sub> cycloalkylalkoxy or cycloalkylalkylthio group optionally substituted by halogen

atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, haloalkyl, alkoxy, haloalkoxy groups;  
a heterocyclyloxy, heterocyclylthio, heterocyclyl-(C<sub>1</sub>-  
C<sub>6</sub>)alkoxy or hetero-cyclyl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio group, option-  
ally substituted by halogen atoms, (C<sub>1</sub>-C<sub>6</sub>) alkyl, haloal-  
5 kyl, alkoxy, halo-alkoxy groups; an aryloxy, arylthio,  
heteroaryloxy, heteroarylthio, aryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryl-  
(C<sub>1</sub>-C<sub>6</sub>)alkylthio, heteroaryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or heteroaryl-  
(C<sub>1</sub>-C<sub>6</sub>)alkylthio group optionally substituted by halogen  
atoms, C<sub>1</sub>-C<sub>6</sub> alkyl groups optionally halogenated, C<sub>1</sub>-C<sub>6</sub>  
10 alkoxy groups optionally halogenated, nitro groups, cyano  
groups;

- A, the same or different when n is greater than or  
equal to 2, represents a halogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl,  
haloalkyl, alkoxy, haloalkoxy group;
- 15 - Y represents an OCH<sub>3</sub> group, an NHCH<sub>3</sub> group, an NH<sub>2</sub>  
group;
- Z represents a CH group or a nitrogen atom N;
- n is an integer ranging from 0 to 4.

Preferred compounds having general formula (I) are  
20 those wherein X<sub>3</sub> represents an R group according to the  
meanings defined above, X<sub>2</sub> and X<sub>4</sub> represent a halogen  
atom, X<sub>1</sub> and X<sub>5</sub> represent a hydrogen atom and n is equal  
to 0.

In the meanings defined above, halogen atom refers  
25 to a fluorine, chlorine, bromine or iodine atom.

Alkyl group refers to a linear or branched group, such as, for example: a methyl, ethyl, iso-propyl, n-propyl, tert-butyl, iso-butyl, n-butyl, n-pentyl, 2,2-dimethylpropyl, n-decyl group.

5        Haloalkyl group refers to a linear or branched group such as, for example: a trifluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, 2,2,2-trifluoro-1-methylethyl group.

Alkoxy or alkylthio group optionally substituted re-  
10    fers to a linear or branched group such as, for example: a methoxy or methylthio group, a difluoromethoxy or difluormethylthio group, a trifluoromethoxy or trifluoromethylthio group, an ethoxy or ethylthio group, a 2,2,2-trifluoroethoxy or 2,2,2-trifluoroethylthio group,  
15    a 2-fluoroethoxy or 2-fluoroethylthio group, a 1,1,2,2-tetrafluoroethoxy or 1,1,2,2-tetrafluoroethylthio group, a 1,2-dichloro-1,2-difluoroethoxy or 1,2-dichloro-1,2-difluoroethylthio group, an n-propoxy or n-propylthio group, an iso-propoxy or iso-propylthio group, a 3-  
20    chloropropoxy or 3-chloropropylthio group, a 3,3,3-trifluoropropoxy or 3,3,3-trifluoropropylthio group, a 1,1,2,3,3,3-hexafluoropropoxy or 1,1,2,3,3,3-hexafluoropropylthio group, an n-butoxy or n-butylthio group, an iso-butoxy or iso-butylthio group, a tert-  
25    butoxy or tert-butylthio group, an n-pentoxy or n-

pentylthio group, a 3-methylbutoxy or 3-methylbutylthio group, a 3,3-dimethylbutoxy or 3,3-dimethylbutylthio, an n-hexyloxy or n-hexylthio group, an n-decyloxy or n-decylthio group, a methoxymethoxy or methoxymethylthio group, an ethoxymethoxy or ethoxymethylthio group, a 2-methoxyethoxy or 2-methoxyethylthio group, a 2-ethoxyethoxy or 2-ethoxyethylthio group, a 2-ethoxypropoxy or 2-ethoxypropylthio group, a 3-methoxyprop-2-yloxy or 3-methoxyprop-2-ylthio group, a 3-ethoxypropoxy or 3-ethoxypropylthio group, a 2-(2-chloroethoxy)ethoxy or 2-(2-chloroethoxy)ethylthio group, a 2-(2-fluoroethoxy)ethoxy or 2-(2-fluoroethoxy)ethylthio group, a 1,1,2-trifluoro-2-trifluoromethoxyethoxy or 1,1,2-trifluoro-2-trifluoromethoxyethylthio group, a 2-(1,1,2,2-tetrafluoroethoxy)ethoxy or 2-(1,1,2,2-tetrafluoroethoxy)ethylthio group, a 2-(2-methoxyethoxy)ethoxy or 2-(2-methoxyethoxy)ethylthio group, a 2-(2-ethoxyethoxy)ethoxy or 2-(2-ethoxyethoxy)ethylthio group, a cyanomethoxy or cyanomethylthio group, a 2-cyanoethoxy or 2-cyanoethylthio group, a 2-aminoethoxy or 2-aminoethylthio group, a 3-aminopropoxy or 3-aminopropylthio group, a 2-(N,N-dimethylamino)ethoxy or 2-(N,N-dimethylamino)ethylthio group, a 2-(N,N-diethylamino)ethoxy or 2-(N,N-diethylamino)ethylthio group, a 3-(N,N-dimethylamino)propoxy or 3-(N,N-dimethylamino)-

propylthio group, a 2-(N,N-diethylamino)propoxy or 2-(N,N-diethylamino)propylthio group, a 2-(N-ethylamino)-ethoxy or 2-(N-ethylamino)ethylthio group, a trimethylsilylmethoxy or trimethylsilylmethylthio group, a 2-(4-chlorophenoxy)ethoxy or 2-(4-chlorophenoxy)-ethylthio group, a 2-(4-methoxyphenoxy)ethoxy or 2-(4-methoxyphenoxy)ethylthio group, a 2-(4-trifluoromethylphenoxy)ethoxy or 2-(4-trifluoromethylphenoxy)-ethylthio group, a 2-(5-trifluoromethyl-2-pyridyloxy)-ethoxy or 2-(5-trifluoromethyl-2-pyridyloxy)ethylthio group, a 3-(5-trifluoromethyl-2-pyridyloxy)propoxy or 3-(5-trifluoromethyl-2-pyridyloxy)propylthio group.

An alkenyloxy or alkenylthio group optionally substituted, refers to a linear or branched group such as, for example: a 2-propenyloxy or 2-propenylthio group, a 3-methyl-2-butenyloxy or 3-methyl-2-butenylthio group, a 3,3-dichloro-2-propenyloxy or 3,3-dichloro-2-propenylthio group, a 3-chloro-4,4,4-trifluoro-2-butenyloxy or 3-chloro-4,4,4-trifluoro-2-butenylthio group, a 3,4,4,4-tetrafluoro-2-butenyloxy or 3,4,4,4-tetrafluoro-2-butenylthio group, a 5,5-dichloro-4-pentenylthio group, a 3-cyano-2-propenyloxy or 3-cyano-2-propenylthio group, a 3-phenyl-2-propenyloxy or 3-phenyl-2-propenylthio group, a 3-(4-chlorophenyl)-2-propenyloxy or 3-(4-chlorophenyl)-2-propenylthio group.

An alkynyloxy or alkynylthio group optionally substituted, refers to a linear or branched group such as, for example, a 2-propynyloxy or 2-propynylthio group, a 3-butyloxy or 3-butylthio group, a 3-butenyl-2-oxy or 3-butenyl-2-thio group, a 3-chloro-2-propynyloxy or 3-chloro-2-propynylthio group, a 4,4,4-trifluoro-2-butyloxy or 4,4,4-trifluoro-2-butylthio group, a 5-chloro-4-pentyloxy or 5-chloro-4-pentylthio group, a 4-methoxy-2-butyloxy or 4-methoxy-2-butylthio group, a 6-methoxy-4-butyloxy or 6-methoxy-4-butylthio group, a 3-(4-chlorophenyl)-2-propynyloxy or 3-(4-chlorophenyl)-2-propynylthio group.

An alkoxyiminoalkylidenoxy or alkoxyiminoalkylidenthio group refers to a linear or branched group such as, for example: a 2-methoxyiminoethylidenoxy or 2-methoxyiminoethylidenthio group, a 2-methoxyimino-propylidenoxy or 2-methoxyiminopropylidenthio group.

Examples of cycloalkoxy or cycloalkylthio groups optionally substituted are: a cyclopentoxy or cyclopentylthio group, a cyclohexyloxy or cyclohexylthio group, a 2,2-difluorocyclohexyloxy or 2,2-difluoro-cyclohexylthio group, a 2,6-dimethylcyclohexyloxy or 2,6-dimethylcyclohexylthio group.

Examples of cycloalkylalkoxy or cycloalkylthio groups optionally substituted are: a cyclopropylmethoxy

or cyclopropylmethylthio group, a 1-cyclopropylethoxy or  
1-cyclopropylethylthio group, a 2,2-dichlorocyclo-  
propylmethoxy or 2,2-dichlorocyclopropylmethylthio group,  
a 2,2-dichloro-1-methylcyclopropylmethoxy or 2,2-  
5 dichloro-1-methylcyclopropylmethylthio group, a 2-  
methylcyclopropylmethoxy or 2-methylcyclopropyl-  
methylthio group, a 2,2-dimethylcyclopropylmethoxy or  
2,2-dimethylcyclopropylmethylthio group, a cyclobutyl-  
methoxy or cyclobutylmethylthio group, a cyclohexyl-  
10 methoxy or cyclohexylmethylthio group.

A heterocyclyl group refers to a mono or polycyclic  
group with 3-14 members, saturated or unsaturated, op-  
tionally benzocondensed but not completely aromatic, con-  
sisting of carbon atoms and one or more heteroatoms, the  
15 same or different, to be selected from nitrogen, sulfur,  
oxygen.

Examples of heterocyclyloxy or heterocyclylthio  
groups optionally substituted, are therefore: a tetrahy-  
drofuranyloxy or tetrahydrofuranylthio group, a tetrahy-  
20 dropyranyloxy or tetrahydropyranylthio group, a tetrahy-  
drothiophenoxy or tetrahydrothiophenylthio group, a 1,3-  
dioxolanyloxy or 1,3-dioxolanylthio group, a 1,4-  
dioxanyloxy or 1,4-dioxanylthio group, a 3-piperidinyloxy  
or 3-piperidinythio group, a 4-piperidinyloxy or 4-  
25 piperidinythio group, a 1-methyl-3-piperidinyloxy or 1-

methyl-3-piperidinylthio group.

Examples of heterocyclyl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or heterocyclyl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio groups optionally substituted, are: a glycidyoxy or glycidylthio group, an oxethanylmethoxy or oxethanymethylthio group, a tetrahydrofuranylmethoxy or tetrahydrofuranylmethylthio group, a tetrahydropyranylmethoxy or tetrahydropyranylmethylthio group, a tetrahydrothiophenylmethoxy or tetrahydrothiophenylmethylthio group, a 1,3-dioxolanylmethoxy or 1,3-dioxolanylmethylthio group, a 1,4-dioxanylmethoxy or 1,4-dioxanymethylthio group, a 1-piperidinylethoxy or 1-piperidinylethylthio group.

An aryl group refers to a mono- or poly-cyclic aromatic carbocyclic group, such as phenyl and naphthyl.

Examples of aryloxy or arylthio groups optionally substituted, are therefore: a phenoxy or phenylthio group, a 4-trifluoromethylphenoxy or 4-trifluoromethylphenylthio group, a 4-chlorophenoxy or 4-chlorophenylthio group, a 4-nitrophenoxy or 4-nitrophenylthio group, a naphthyloxy or naphthylthio group.

A hetero-aryl group refers to a mono or polycyclic aromatic group consisting of carbon atoms and one or more hetero-atoms, the same or different, to be selected from nitrogen, sulfur, oxygen; for example: pyridyl, thio-

phenyl, furanyl, pyrrolyl, pyrazolyl, pyrimidyl, tri-  
azolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl,  
thiadiazolyl, triazinyl, tetrazinyl, benzoxazolyl.

Examples of hetero-aryloxy or hetero-arylthio groups  
5 optionally substituted are therefore: 2-pyridyloxy or 2-  
pyridylthio, 5-chloro-2-pyridyloxy or 5-chloro-2-  
pyridylthio, 5-trifluoromethyl-2-pyridyloxy or 5-  
trifluoromethyl-2-pyridylthio, 5-trifluoromethyl-1,3,4-  
thiadiazolyloxy or 5-trifluoromethyl-1,3,4-thiadiazol-  
10 ylthio.

Examples of aryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or aryl-(C<sub>1</sub>-  
C<sub>6</sub>)alkylthio groups optionally substituted, are: a benzy-  
loxy or benzylthio group, a 4-chlorobenzoyloxy or 4-  
chlorobenzylthio group, a 4-tert-butylbenzyloxy or 4-  
15 tert-butylbenzylthio group, a 1-(4-chlorophenyl)ethoxy or  
1-(4-chlorophenyl)ethylthio group.

Examples of hetero-aryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or hetero-  
aryl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio groups optionally substituted, are:  
2-furanylmethoxy or 2-furanylmethylthio, 5-nitro-2-  
20 furanylmethoxy or 5-nitro-2-furanylmethylthio, 2-  
thienylmethoxy or 2-thienylmethylthio, 3-pyridylmethoxy  
or 3-pyridylmethylthio, 6-chloro-3-pyridylmethoxy or 6-  
chloro-3-pyridylmethylthio.


Although the compounds having general formula (I)  
25 partially fall within the scope of the general formulae

of patent applications EP 178826, EP 226917, EP 252213, EP 278595 and EP 398692, they are not specifically illustrated therein and are consequently new.

Compounds having formula (I) which are interesting for their biological activity are those described in Examples 1-4 provided further on.

A further object of the present invention relates to the process for the preparation of the compounds having general formula (I).

The compounds having general formula (I) can be obtained by the condensation of a compound having general formula (II) with a phenol having general formula (III), according to reaction scheme 1:



in said formulae,  $X_1$ ,  $X_2$ ,  $X_3$ ,  $X_4$ ,  $X_5$ ,  $A$ ,  $Y$ ,  $Z$  and  $n$  have the meanings defined above,  $L$  represents a leaving group such as a chlorine atom, a bromine atom or a  $R_L\text{SO}_3^-$  group wherein  $R_L$  represents a  $C_1$ - $C_6$  alkyl or haloalkyl, or a

phenyl optionally substituted.

The reaction can be conveniently carried out in an inert organic solvent, at a temperature ranging from 0°C to the boiling point of the reaction mixture, optionally  
5 in the presence of an inorganic or organic base.

Preferred solvents for the reaction are alcohols (methanol, ethanol, propanol, methoxyethanol, etc.), ethers (ethyl ether, isopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane, etc.), esters (ethyl acetate,  
10 etc.), ketones (acetone, methylethylketone, etc.), chlorinated hydrocarbons (methylene chloride, dichloroethane, chloroform, carbon tetrachloride, etc.), aromatic hydrocarbons (benzene, toluene, xylene, etc.), aliphatic hydrocarbons (hexane, heptane, cyclohexane, etc.), aprotic  
15 dipolar solvents (N,N-dimethylformamide, dimethylsulfoxide, sulfolane, etc.).

Preferred inorganic bases are: hydrides, hydroxides, carbonates of alkaline or alkaline-earth metals (sodium, potassium, calcium, etc.).

20 Preferred organic bases are: pyridine, dimethylaminopyridine, aliphatic amines (diethylamine, triethylamine, etc.), cyclic amines (morpholine, piperidine, diazabicycloundecene, etc.), alcoholates of alkaline metals (sodium methylate, potassium t-butyrate, etc.).

25 The intermediates having general formula (II) are

known compounds.

The phenols of general formula (III), when not known in themselves, can be prepared according to methods known in organic chemical practice.

5       The compounds having general formula (I) wherein Y represents  $\text{NH}_2$  or  $\text{NHCH}_3$  can be alternatively prepared by reaction of the corresponding methyl ester [ $\text{Y} = \text{OCH}_3$  in general formula (I)] with ammonia ( $\text{NH}_3$ ), or with methylamine ( $\text{CH}_3\text{NH}_2$ ).

10       The reaction can be conveniently carried out in an inert organic solvent, at a temperature ranging from  $0^\circ\text{C}$  to the boiling point of the reaction mixture.

Preferred solvents for carrying out the reaction are alcohols (methanol, ethanol, propanol, methoxyethanol, etc.), ethers (ethyl ether, isopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane, etc.), ketones (acetone, methylethylketone, etc.), chlorinated hydrocarbons (methylene chloride, dichloroethane, chloroform, carbon tetrachloride, etc.), aromatic hydrocarbons (benzene, toluene, xylene, etc.), aliphatic hydrocarbons (hexane, heptane, cyclohexane, etc.), aprotic dipolar solvents (N,N-dimethylformamide, dimethylsulfoxide, sulfolane, etc.) or mixtures of these in any ratio.

25       The compounds having general formula (I) have an geometric isomerism on the  $\text{C}=\text{Z}$  double bond; the use of

the compounds having formula (I) as isomeric mixtures in any proportion, as also the production and use of the single E or Z isomers are included in the scope of the present invention. For the purposes of a biological activity E isomers of the compounds of formula (I) are preferred.

The compounds having general formula (I) have a high acaricidal and insecticidal activity which is exerted with respect to the adults, larvae and eggs of mites and insects which are harmful in the agrarian, civil and zoo-technical field.

A further object of the present invention therefore relates to the use of the compounds having general formula (I) as acaricides and/or insecticides and/or fungicides, both in agriculture and other fields. In particular, the use of E isomers of the compounds having formula (I) are preferred.

The use of the compounds having formula (I) wherein  $X_3$  represents an R group according to the meanings defined above,  $X_2$  and  $X_4$  represent a halogen atom,  $X_1$  and  $X_5$  represent a hydrogen atom and n is equal to 0, is preferred.

Particularly preferred is the use of compounds having formula (I) selected from:

- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophen-

- oxymethyl)phenyl]-3-methoxyacrylate;
- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-2-methoxyiminoacetate;
- (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy-  
5 methyl)phenyl]-N-methyl-2-methoxyiminoacetamide;
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]phenyl}-2-methoxyiminoacetate;
- 10 - (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]phenyl}-3-methoxyacrylate;
- 15 - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]phenyl}-2-methoxyiminoacetate;
- (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide;
- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxymethyl]phenyl}-3-methoxy-  
20 acrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxymethyl]phenyl}-2-methoxyiminoacetate;
- 25 - (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-

2-enyloxy)phenoxyethyl]phenyl}-N-methyl-2-methoxyimino-acetamide;

- methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichlorophenoxyethyl]phenyl]-3-methoxyacrylate;

5 - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy)phenoxyethyl]phenyl}-3-methoxyacrylate;

- methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy)phenoxyethyl]phenyl}-3-methoxyacrylate;

- methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichlorophenoxyethyl]phenyl]-3-methoxyacrylate;

- methyl (E)-2-{2-[3,5-dichloro-4-(2,4-dichlorobenzyloxy)phenoxyethyl]phenyl}-3-methoxyacrylate;

- methyl (E)-2-{2-[3,5-dichloro-4-(4-chlorobenzyloxy)phenoxyethyl]phenyl}-3-methoxyacrylate.

15 In particular, the compounds having general formula (I) are active against important species of tetranychidae (*Tetranychus urticae*, *Tetranychus telarius*, *Tetranychus cinnabarinus*, *Eotetranychus carpini*, *Panonychus ulmi*, *Panonychus citri*, etc.), eriophyidae (*Phytoptus avel-*  
20 *lanae*, *Eriophyes vitis*, *Eriophyes piri*, etc.) tarsonemidae (*Steneotarsonemus pallidus*, etc.), hemiptera (*Macrosiphum euphorbiae*, *Aphis fabae*, *Myzus persicae*, etc.), lepidoptera (*Spodoptera* spp., *Heliothis* spp., *Chilo* spp., *Carpocapsa pomonella*, etc.), coleoptera (*Leptinotarsa de-*  
25 *cemlineata*, *Phaedon cochleariae*, etc.), diptera (*Aedes*

spp., *Culex* spp., *Musca* spp., etc.).

The compounds having general formula (I) also have a good fungicidal activity of both a preventive and curative nature: examples of phytopathogen fungi which can be controlled with the compounds of the invention are: Helminthosporium spp., Erysiphe spp., Puccinia spp., *Plasmodiophora viticola*, Pythium spp., Phytophthora spp., Rhynchosporium spp., Septoria spp., *Sphaerotheca fuliginea*, *Podosphaera leucotricha*, *Pyricularia oryzae*, *Uncinula necator*, Venturia spp., *Botrytis cinerea*, Fusarium spp., Alternaria spp., Cercospora spp.

At the same time, the compounds having general formula (I) have a low toxicity with respect to many useful insects and mites, with respect to mammals, fish, birds, and have no phytotoxicity.

Thanks to their positive characteristics, they can be advantageously used in defending not only crops of agrarian and horticultural interest, but also domestic and breeding animals, as well as environments frequented by human beings, from harmful mites, insects and fungi.

The quantity of compound to be applied to obtain the desired effect can vary in relation to various factors such as, for example, the compound used, the crop to be preserved, the type of harmful organism, the degree of infestation, the climatic conditions, the method of ap-

plication, the formulation adopted.

Doses of compound ranging from 10 g to 5 kg per hectare generally provide a sufficient control.

A further object of the present invention also relates to a method for controlling mites and/or insects and/or phytopathogenic fungi in crops of agrarian and horticultural interest and/or on domestic and breeding animals and/or in environments frequented by human beings, by the application of the compounds having general formula (I). In particular, the quantity of compound to be applied varies from 10 g to 5 kg per hectare.

For practical use in agriculture, it is often useful to use compositions containing one or more compounds having general formula (I).

A further object of the present invention therefore relates to acaricidal and/or insecticidal and/or fungicidal compositions containing one or more compounds having general formula (I) as active principle.

Compositions can be used in the form of dry powders, wettable powders, emulsifiable concentrates, micro-emulsions, pastes, granulates, solutions, suspensions, etc.: the selection of the type of composition depends on the specific use.

The compositions are prepared in the known way, for example by diluting or dissolving the active substance

with a solvent medium and/or a solid diluent, optionally in the presence of surface-active agents.

Solid diluents, or carriers which can be used are, for example: silica, kaolin, bentonite, talc, diatomite, dolomite, calcium carbonate, magnesia, chalk, clays, synthetic silicates, attapulgite, sepiolite.

Liquid diluents which can be used are, for example, in addition to water, aromatic organic solvents (xvols or mixtures of alkylbenzols, chlorobenzene, etc.), paraffins (petroleum fractions), alcohols (methanol, propanol, butanol, octanol, glycerin, etc.), esters (ethyl acetate, isobutyl acetate, etc.), ketones (cyclohexanone, acetone, acetophenone, isophorone, ethylamylketone, etc.), amides (N,N-dimethylformamide, N-methylpyrrolidone, etc.).

Surface-active agents which can be used are salts of sodium, calcium, triethylamine or triethanolamine of alkylsulfonates, alkylarylsulfonates, polyethoxylated alkylphenols, polyethoxylated esters of sorbitol, lignin-sulfonates, etc.

The compositions can also contain special additives for particular purposes, for example adhesion agents such as Arabic gum, polyvinyl alcohol, polyvinyl-pyrrolidone, etc.

The concentration of active principle in the above compositions can vary within a wide range depending on

the active compound, the applications for which they are destined, the environmental conditions and the type of formulation adopted. In general the concentration of active principle ranges from 1 to 90%, preferably from 5 to 50%.

If required, it is possible to add to the compositions, other active principles compatible with the compounds having general formula (I), such as, for example, other acaricides/insecticides, fungicides, phyto-  
10 regulators, antibiotics, herbicides, fertilizers.

Examples of other acaricides/insecticides which can be added to the above compositions are: abamectin, acephate, acetamiprid, acetoprole, acrinathrin, acequinocyl, alanycarb, aldicarb, allethrin, alpha-cypermethrin, 15 amitraz, azadirachtin, azamethiphos, azinphos-ethyl, azinphos-methyl, azocyclotin, bendiocarb, benfuracarb, bensultap, benzoximate, bifenazate, bifenthrin, bioallethrin, bioresmethrin, bistrifluron, (DBI-3204), bromopropylate, buprofezin, butocarboxim, butoxycarboxim, cadusafos, 20 carbofuran, carbosulfan, cartap, CGA 50439, chinomethionat, chlordane, chlorethoxyfos, chlorfenapyr, chlorfenvinphos, chlorfluazuron, chlormephos, chloropicrin, chlorpyrifos, chlorpyrifos-methyl, chromafenozide, clofentezine, clothianidin, coumaphos, cyanophos, cycloprothrin, 25 cyfluthrin, cyhalothrin, cyhexatin, cypermethrin,

cyphenothrin, cyromazine, DDT, deltamethrin, demeton-S-methyl, diafenthiuron, diazinon, dichlorvos, dicofol, dicrotophos, dicyclanil, dienochlor, diflubenzuron, dimethoate, dimethylvinphos, dinobuton, dinocap, dinotefuran (MTI-446), diofenolan, disulfoton, DNOC, indoxacarb, endosulfan, EPN, esfenvalerate, ethiofencarb, ethion, ethiprole, ethoprophos, etofenprox, etoxazole, etrimfos, famphur, fenazaquin, fenbutatin oxide, fenitrothion, fenobucarb, fenothiocarb, fenoxycarb, fenpropathrin, fenpyroximate, fenthion, fenvalerate, fipronil, flucycloxuron, flucythrinate, flufenoxuron, flufenprox, flufenzin, flumethrin, fluvalinate, fonofos, formetanate, formothion, furathiocarb, halfenprox, halofenozide, heptachlor, heptenophos, hexaflumuron, hexythiazox, hydroprene, IKA-2000, IKI-220, imidacloprid, isazofos, isofenphos, isoprocarb, isoxathion, lindane, lufenuron, malathion, mecarbam, methacrifos, methamidophos, methidathion, methiocarb, methomyl, methoprene, methoxychlor, methoxyfenozide, metolcarb, mevinphos, milbemectin, monocrotophos, naled, nicotine, nitenpyran, nithiazine, novaluron, omethoate, oxamyl, oxydemeton-methyl, parathion, permethrin, petroleum oils, phenothrin, phenthoate, phorate, phosalone, phosmet, phosphamidon, phoxim, pirimicarb, pirimiphos-ethyl, pirimiphos-methyl, profenofos, propaphos, propargite, propetamphos, propoxur, prothiofos,

protrifenbute, pymetrozine, pyraclofos, pyrethrins,  
pyridaben, pyridaphenthion, pyrimidifen, pyriproxyfen,  
quinalphos, rotenone, S-1812, silafluofen, spirodiclofen  
(BAJ2740), sulfluramid, sulfotep, sulprofos, tar oils,  
5 tebufenozide, tebufenpyrad, tebupirimfos, teflubenzuron,  
tefluthrin, temephos, terbufos, tetrachlorvinphos, tetra-  
difon, tetramethrin, thiacloprid, thiamethoxam, thiocy-  
clam, thiodicarb, thiofanox, thiometon, tolfenpyrad, tra-  
lomethrin, transfluthrin, triazamate, triazophos, tri-  
10 chlorfon, triflumuron, trimethacarb, vamidothion, spino-  
sad, vaniliprole, XMC, xylylcarb, ZXI 8901.

Some examples are provided for illustrative pur-  
poses but in no way limit the scope of the present inven-  
tion.

15

#### EXAMPLE 1

Preparation of methyl (E)-2-[2-(4-cyclopropylmethoxy-  
3,5-dichlorophenoxy)methyl]phenyl]-3-methoxyacrylate (com-  
pound Nr. 1).

20

A solution of 4-cyclopropylmethoxy-3,5-dichloro-  
phenol (1.54 g; 6.6 mmol) in DMF (5 ml) is added drop-  
wise, at 0°C, to a suspension of sodium hydride (0.16 g;  
6.6 mmol) in N,N-dimethyl formamide (DMF; 5 ml). The  
mixture is kept under stirring at room temperature for 30  
25 minutes and a solution of methyl (E)-2-(2-

bromomethylphenyl)-3-methoxyacrylate (1.9 g; 6.6 mmoles) in DMF (5 ml) is then added. The mixture is kept under stirring for 4 hours; the mixture is diluted with water (40 ml) and an extraction is effected with ethyl acetate (3x25 ml). The organic phases are joined, washed with water (2x20 ml), dried with sodium sulfate, filtered and concentrated at reduced pressure. The raw product obtained is purified by means of silica gel chromatography, eluting with hexane/ethyl acetate 9:1. 1.9 g of the desired product (thick oil) are obtained.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.33 (2H,m), 0.61 (2H,m), 1.30 (1H,m), 3.71 (3H,s), 3.78 (2H,d), 3.85 (3H,s), 4.90 (2H,s), 6.82 (2H,s), 7.10-7.55 (4H,m), 7,61 (1H,s).

#### EXAMPLE 2

15 Preparation of methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy)methyl)-phenyl]-2-methoxyiminoacetate (compound Nr.2).

A solution of 4-cyclopropylmethoxy-3,5-dichlorophenol (1.54 g; 6.6 mmoles) in DMF (5 ml) is added dropwise, at 0°C, to a suspension of sodium hydride (0.16 g; 6.6 mmoles) in DMF (5 ml). The mixture is kept under stirring at room temperature for 30 minutes, and a solution of methyl (E)-2-(2-bromomethylphenyl)-3-methoxyiminoacetate (1.9 g; 6.6 mmoles) in DMF (5 ml) is then added. The mixture is kept under stirring for 4

hours; the mixture is diluted with water (40 ml) and an extraction is effected with ethyl acetate (3x25 ml). The organic phases are joined, washed with water (2x20 ml), dried with sodium sulfate, filtered and concentrated at reduced pressure. The raw product obtained is purified by means of silica gel chromatography, eluting with hexane/ethyl acetate 9:1. 1.7 g of the desired product (thick oil) are obtained.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.33 (2H,m), 0.61 (2H,m), 1.30 (1H,m), 3.78 (2H,d), 3.85 (3H,s), 4.03 (3H,s), 4.90 (2H,s), 6.82 (2H,s), 7.10-7.55 (4H,m).

### EXAMPLE 3

Preparation of (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy)methyl]-phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.3).

A 40% aqueous solution of methylamine (0.8 ml) is added to a solution of methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy)methyl]-phenyl]-2-methoxyiminoacetate (compound nr.2; 0.88g; 2 mmols) in DMF (2 ml) and methanol (4 ml). The mixture is kept under stirring for 2 hours at room temperature, diluted with water (40 ml) and an extraction is effected with ethyl acetate (3x25 ml). The organic phases are joined, washed with water (2x20 ml), dried with sodium sulfate and concentrated at reduced pressure. 0.8 g of the product

(thick oil) are obtained, which do not require further purification.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.33 (2H,m), 0.61 (2H,m), 1.30 (1H,m),  
2.90 (3H,d), 3.78 (2H,d), 4.03 (3H,s), 4.90 (2H,s), 6.78  
5 (1H,mb), 6.82 (2H,s), 7.10-7.55 (4H,m).

#### EXAMPLE 4

Preparation of compounds Nr. 4-146.

Operating as described in examples 1-3, the following products were prepared and identified by means of <sup>1</sup>H-NMR spectroscopy and GC-MS:

- methyl (E)-2-[2-(4-cyclopentoxy-3,5-dichlorophenoxy-methyl)phenyl]-3-methoxyacrylate (compound Nr.4);
- methyl (E)-2-[2-(4-cyclopentoxy-3,5-dichlorophenoxy-methyl)phenyl]-2-methoxyiminoacetate (compound Nr.5);
- 15 - (E)-2-[2-(4-cyclopentoxy-3,5-dichlorophenoxy-methyl)phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.6);
- methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichlorophenoxy-methyl)phenyl]-3-methoxyacrylate (compound Nr.7);
- methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichlorophenoxy-methyl)phenyl]-2-methoxyiminoacetate (compound  
20 Nr.8);
- (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichlorophenoxy-methyl)phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.9);
- 25 - methyl (E)-2-[2-(4-cyclohexyloxy-3,5-dichlorophe-

- nooxymethyl)phenyl]-3-methoxyacrylate (compound Nr.10);
- methyl (E)-2-[2-(4-cyclohexyloxy-3,5-dichlorophenoxy)methyl)phenyl]-2-methoxyiminoacetate (compound Nr.11);
  - (E)-2-[2-(4-cyclohexyloxy-3,5-dichlorophenoxy)methyl)phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.12);
  - methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichlorophenoxy)methyl)phenyl]-3-methoxyacrylate, solid with m.p. 103°C;
- 10 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.22 (5H,m), 1.88 (6H,m), 3.73 (3H,s), 3.74 (2H,d), 3.82 (3H,s), 4.93 (2H,s), 6.84 (2H,s), 7.10-7.55 (4H,m), 7.62 (1H,s). (compound Nr.13);
- methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichlorophenoxy)methyl)phenyl]-2-methoxyiminoacetate (compound Nr.14);
- 15 - (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichlorophenoxy)methyl)phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.15);
- methyl (E)-2-[2-(3,5-dichloro-4-methoxyphenoxy)methyl)phenyl]-3-methoxyacrylate (compound Nr.16);
- 20 - methyl (E)-2-[2-(3,5-dichloro-4-methoxyphenoxy)methyl)phenyl]-2-methoxyiminoacetate (compound Nr.17);
- (E)-2-[2-(3,5-dichloro-4-methoxyphenoxy)methyl)phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.18);
  - methyl (E)-2-[2-(3,5-dichloro-4-ethoxyphenoxy)methyl)phenyl]-3-methoxyacrylate (compound Nr.19);
- 25

- methyl (E)-2-[2-(3,5-dichloro-4-ethoxyphenoxy)methyl]phenyl]-2-methoxyiminoacetate (compound Nr.20);
- (E)-2-[2-(3,5-dichloro-4-ethoxyphenoxy)methyl]phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.21);
- 5 - methyl (E)-2-[2-(3,5-dichloro-4-isopropoxyphenoxy)methyl]phenyl]-3-methoxyacrylate (compound Nr.22);
- methyl (E)-2-[2-(3,5-dichloro-4-isopropoxyphenoxy)methyl]phenyl]-2-methoxyiminoacetate (compound Nr.23);
- (E)-2-[2-(3,5-dichloro-4-isopropoxyphenoxy)methyl]phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.24);
- 10 - methyl (E)-2-[2-(3,5-dichloro-4-isobutoxyphenoxy)methyl]phenyl]-3-methoxyacrylate (compound Nr.25);
- methyl (E)-2-[2-(3,5-dichloro-4-isobutoxyphenoxy)methyl]phenyl]-2-methoxyiminoacetate (compound Nr.26);
- 15 - (E)-2-[2-(3,5-dichloro-4-isobutoxyphenoxy)methyl]phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.27);
- methyl (E)-2-{2-[3,5-dichloro-4-(2,2-dimethylpropoxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.28);
- 20 - methyl (E)-2-{2-[3,5-dichloro-4-(2,2-dimethylpropoxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate (compound Nr.29);
- (E)-2-{2-[3,5-dichloro-4-(2,2-dimethylpropoxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.30);
- 25

- methyl (E)-2-{2-[3,5-dichloro-4-(2,2-dimethylethoxy)phoxymethyl]phenyl}-3-methoxyacrylate (compound Nr.31);
- methyl (E)-2-{2-[3,5-dichloro-4-(2,2-dimethylethoxy)phoxymethyl]phenyl}-2-methoxyiminoacetate (compound  
5 Nr.32);
- (E)-2-{2-[3,5-dichloro-4-(2,2-dimethylethoxy)phoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.33);
- methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy)phoxymethyl]phenyl}-3-methoxyacrylate, m.p. 70°C;  
10 <sup>1</sup>H-NMR(CDCl<sub>3</sub>): δ 1.0 (6H,d), 1.73 (2H,m), 1.94 (1H,m), 3.73 (3H,s), 3.82 (3H,s), 3.98 (2H,m), 4.93 (2H,s), 6.84 (2H,s), 7.10-7.55 (4H,m), 7.62 (1H,s). (compound Nr.34);
- methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy)phoxymethyl]phenyl}-2-methoxyiminoacetate (compound  
15 Nr.35);
- (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy)phoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.36);
- 20 - methyl (E)-2-[2-(3,5-dichloro-4-hexyloxyphoxymethyl)phenyl]-3-methoxyacrylate (compound Nr.37);
- methyl (E)-2-[2-(3,5-dichloro-4-hexyloxyphoxymethyl)phenyl]-2-methoxyiminoacetate (compound Nr.38);
- (E)-2-[2-(3,5-dichloro-4-hexyloxyphoxymethyl)  
25 phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.39);

- methyl (E)-2-[2-(3,5-dichloro-4-decyloxyphenoxy-methyl)phenyl]-3-methoxyacrylate (compound Nr.40);
- methyl (E)-2-[2-(3,5-dichloro-4-decyloxyphenoxy-methyl)phenyl]-2-methoxyiminoacetate (compound Nr.41);
- 5 - (E)-2-[2-(3,5-dichloro-4-decyloxyphenoxy-methyl)-phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.42);
- methyl (E)-2-[2-(3,5-dichloro-4-n-propoxyphenoxy-methyl)phenyl]-3-methoxyacrylate (compound Nr.43);
- methyl (E)-2-[2-(3,5-dichloro-4-n-propoxyphenoxy-methyl)phenyl]-2-methoxyiminoacetate (compound Nr.44);
- 10 - (E)-2-[2-(3,5-dichloro-4-n-propoxyphenoxy-methyl)-phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.45);
- methyl (E)-2-{2-[3,5-dichloro-4-(2-ethoxyethoxy)-phenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.46);
- 15 - methyl (E)-2-{2-[3,5-dichloro-4-(2-ethoxyethoxy)-phenoxy-methyl]phenyl}-2-methoxyaminoacetate (compound Nr.47);
- (E)-2-{2-[3,5-dichloro-4-(2-ethoxyethoxy)phenoxy-methyl]phenyl}-2-methoxyaminoacetamide (compound Nr.48);
- 20 - methyl (E)-2-{2-[3,5-dichloro-4-(2-methoxyethoxy)-phenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.49);
- methyl (E)-2-{2-[3,5-dichloro-4-(2-methoxyethoxy)-phenoxy-methyl]phenyl}-2-methoxyaminoacetate (compound Nr.50);
- 25 - (E)-2-{2-[3,5-dichloro-4-(2-methoxyethoxy)phenoxy-methyl]phenyl}-2-methoxyaminoacetamide (compound Nr.51);

- thyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.51);
- methyl (E)-2-{2-[3,5-dichloro-4-(2-ethoxymethoxy)-phoxymethyl]phenyl}-3-methoxyacrylate (compound Nr.52);
- 5 - methyl (E)-2-{2-[3,5-dichloro-4-(2-ethoxymethoxy)-phoxymethyl]phenyl}-2-methoxyiminoacetate (compound Nr.53);
- (E)-2-{2-[3,5-dichloro-4-(2-ethoxymethoxy)-phoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound
- 10 Nr.54);
- methyl (E)-2-{2-[3,5-dichloro-4-(2,2,2-trifluoroethoxy)phoxymethyl]phenyl}-3-methoxyacrylate (compound Nr.55);
- methyl (E)-2-{2-[3,5-dichloro-4-(2,2,2-trifluoroethoxy)phoxymethyl]phenyl}-2-methoxyiminoacetate (com-
- 15 pound Nr.56);
- (E)-2-{2-[3,5-dichloro-4-(2,2,2-trifluoroethoxy)-phoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.57);
- 20 - methyl (E)-2-[2-(3,5-dichloro-4-trifluoromethoxy-phoxymethyl)phenyl]-3-methoxyacrylate (compound Nr.58);
- methyl (E)-2-[2-(3,5-dichloro-4-trifluoromethoxy-phoxymethyl)phenyl]-2-methoxyiminoacetate (compound
- Nr.59);
- 25 - (E)-2-[2-(3,5-dichloro-4-trifluoromethoxyphoxy-

methyl]phenyl]-N-methyl-2-methoxyiminoacetamide (compound Nr.60);

5 - methyl(E)-2-{2-[3,5-dichloro-4-(1,1,2,2-tetrafluoroethoxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.61);

- methyl(E)-2-{2-[3,5-dichloro-4-(1,1,2,2-tetrafluoroethoxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate (compound Nr.62);

10 - (E)-2-{2-[3,5-dichloro-4-(1,1,2,2-tetrafluoroethoxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.63);

- methyl(E)-2-{2-[3,5-dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.64);

15 - methyl(E)-2-{2-[3,5-dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate (compound Nr.65);

- (E)-2-{2-[3,5-dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.66);

- methyl(E)-2-{2-[3,5-dichloro-4-(2-trifluoromethoxy-1,1,2-trifluoroethoxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.67);

25 - methyl(E)-2-{2-[3,5-dichloro-4-(2-trifluoromethoxy-1,1,2-trifluoroethoxy)phenoxy)methyl]phenyl}-2-methoxy-

iminoacetate (compound Nr.68);

- (E)-2-{2-[3,5-dichloro-4-(2-trifluoromethoxy-1,1,2-trifluoroethoxy)phenoxyethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.69);

5 - methyl (E)-2-{2-[3,5-dichloro-4-(2-(2-ethoxyethoxy)-ethoxy)phenoxyethyl]phenyl}-3-methoxyacrylate (compound Nr.70);

- methyl (E)-2-{2-[3,5-dichloro-4-(2-(2-ethoxyethoxy)-ethoxy)phenoxyethyl]phenyl}-3-methoxyiminoacetate (compound Nr.71);

- (E)-2-{2-[3,5-dichloro-4-(2-(2-ethoxyethoxy)ethoxy)-phenoxyethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.72);

- methyl (E)-2-{2-[3,5-dichloro-4-(4-trifluoromethylphenoxy)phenoxyethyl]phenyl}-3-methoxyacrylate (compound Nr.73);

- methyl (E)-2-{2-[3,5-dichloro-4-(4-trifluoromethylphenoxy)phenoxyethyl]phenyl}-2-methoxyiminoacetate (compound Nr.74);

20 - (E)-2-{2-[3,5-dichloro-4-(4-trifluoromethylphenoxy)-phenoxyethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.75);

- methyl (E)-2-{2-[3,5-dichloro-4-(4-cyanophenoxy)-phenoxyethyl]phenyl}-3-methoxyacrylate (compound Nr.76);

25 - methyl (E)-2-{2-[3,5-dichloro-4-(4-cyanophenoxy)-

- phenoxymethyl]phenyl}-2-methoxyiminoacetate (compound  
Nr.77);
- (E)-2-{2-[3,5-dichloro-4-(4-cyanophenoxy)phenoxy-  
methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound  
5 Nr.78);
- methyl (E)-2-{2-[3,5-dichloro-4-(4-nitrophenoxy)-  
phenoxymethyl]phenyl}-3-methoxyacrylate (compound Nr.79);
- methyl (E)-2-{2-[3,5-dichloro-4-(4-nitrophenoxy)-  
phenoxymethyl]phenyl}-2-methoxyiminoacetate (compound  
10 Nr.80);
- (E)-2-{2-[3,5-dichloro-4-(4-nitrophenoxy)phenoxy-  
methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound  
Nr.81);
- methyl (E)-2-[2-(3,5-dichloro-4-benzyloxyphenoxy-  
15 methyl)phenyl]-3-methoxyacrylate (compound Nr.82);
- methyl (E)-2-[2-(3,5-dichloro-4-benzyloxyphenoxy-  
methyl)phenyl]-2-methoxyiminoacetate (compound Nr.83);
- (E)-2-[2-(3,5-dichloro-4-benzyloxyphenoxy-methyl)-  
phenyl]-N-methyl-2-methoxyiminoacetamide (compound  
20 nr.84);
- methyl (E)-2-{2-[3,5-dichloro-4-(4-methoxybenzyloxy)  
phenoxymethyl]phenyl}-3-methoxyacrylate (compound Nr.85);
- methyl (E)-2-{2-[3,5-dichloro-4-(4-methoxybenzyloxy)  
phenoxymethyl]phenyl}-2-methoxyiminoacetate (compound  
25 Nr.86);

- (E)-2-{2-[3,5-dichloro-4-(4-methoxybenzyloxy)-phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.87);
- methyl (E)-2-{2-[3,5-dichloro-4-(4-tert-butyl)benzyloxy)phenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.88);
- methyl (E)-2-{2-[3,5-dichloro-4-(4-tert-butyl) benzyloxy)phenoxy-methyl]phenyl}-2-methoxyiminoacetate (compound Nr.89);
- 10 - (E)-2-{2-[3,5-dichloro-4-(4-tert-butyl)benzyloxy)phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.90);
- methyl (E)-2-{2-[3,5-dichloro-4-(5-trifluoromethylpyrid-2-yloxy)phenoxy-methyl]phenyl}-3-methoxyacrylate
- 15 (compound Nr.91);
- methyl (E)-2-{2-[3,5-dichloro-4-(5-trifluoromethylpyrid-2-yloxy)phenoxy-methyl]phenyl}-2-methoxyiminoacetate (compound Nr.92);
- (E)-2-{2-[3,5-dichloro-4-(5-trifluoromethylpyrid-2-yloxy)phenoxy-methyl]phenyl}-N-methyl-2-methoxyimino-
- 20 acetamide (compound Nr.93);
- methyl (E)-2-{2-[3,5-dichloro-4-(2-propynyloxy)-phenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.94);
- methyl (E)-2-{2-[3,5-dichloro-4-(2-propynyloxy)phenoxy-methyl]phenyl}-2-methoxyiminoacetate (compound Nr.95);
- 25

- (E)-2-{2-[3,5-dichloro-4-(2-propynyloxy)-phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.96);
- methyl (E)-2-{2-[3,5-dichloro-4-(3-butyne-2-yloxy)-  
5 phenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.97);
- methyl (E)-2-{2-[3,5-dichloro-4-(3-butyne-2-yloxy)-phenoxy-methyl]phenyl}-2-methoxyiminoacetate (compound Nr.98);
- (E)-2-{2-[3,5-dichloro-4-(3-butyne-2-yloxy)phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound  
10 Nr.99);
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxy-methyl]phenyl}-3-methoxyacrylate, thick oil;
- 15  $^1\text{H-NMR}(\text{CDCl}_3)$ :  $\delta$  3.74 (3H,s), 3.82 (3H,s), 4.60 (2H,d), 4.93 (2H,s), 6.30 (1H,t), 6.84 (2H,s), 7.10-7.55 (4H,m), 7.62 (1H,s). (compound Nr.100);
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxy-methyl]phenyl}-2-methoxyiminoacetate (com-  
20 pound Nr.101);
- (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.102);
- methyl (E)-2-{2-[3,5-dichloro-4-(cyanomethoxy)pheno-  
25 xymethyl]phenyl}-3-methoxyacrylate (compound Nr.103);

- methyl (E)-2-{2-[3,5-dichloro-4-(cyanomethoxy)phenoxy-methyl]phenyl}-2-methoxyiminoacetate (compound Nr.104);
- (E)-2-{2-[3,5-dichloro-4-(cyanomethoxy)phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound  
5 Nr.105);
- methyl (E)-2-{2-[2,6-dichloro-4-(2,2-dimethylethyl)phenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.106);
- methyl (E)-2-{2-[2,6-dichloro-4-(2,2-dimethylethyl)phenoxy-methyl]phenyl}-2-methoxyiminoacetate (compound  
10 Nr.107);
- (E)-2-{2-[2,6-dichloro-4-(2,2-dimethylethyl)phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.108);
- methyl (E)-2-[2-(3-cyclopropylmethoxy-4,6-dichloro-  
15 phenoxy-methyl)phenyl]-3-methoxyacrylate (compound Nr.109);
- methyl (E)-2-[2-(3-cyclopropylmethoxy-4,6-dichloro-phenoxy-methyl)phenyl]-2-methoxyiminoacetate (compound Nr.110);
- (E)-2-[2-(3-cyclopropylmethoxy-4,6-dichlorophenoxy-methyl)phenyl]-N-methyl-2-methoxyiminoacetamide (compound  
20 Nr.111);
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxy-methyl]phenyl}-3-methoxyacrylate (compound Nr.112);
- 25 - methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-

3,5-dichlorophenoxymethyl]phenyl}-2-methoxyiminoacetate  
(compound Nr.113);

- (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-  
dichlorophenoxymethyl]phenyl}-N-methyl-2-methoxyimino-  
5 acetamide (compound Nr.114);

- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-  
trifluorobut-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxy-  
acrylate, thick oil,

<sup>1</sup>H-NMR(CDCl<sub>3</sub>): δ 3.74 (3H,s), 3.82 (3H,s), 4.72 (2H,m),  
10 4.93 (2H,s), 6.72 (1H,m), 6.84 (2H,s), 7.10-7.55 (4H,m),  
7.62 (1H,s). (compound Nr.115);

- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-  
trifluorobut-2-enyloxy)phenoxy)methyl]phenyl}-2-methoxy-  
iminoacetate (compound Nr.116);

15 - (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluoro-  
but-2-enyloxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxy-  
iminoacetamide (compound Nr.117);

- methyl (E)-2-{2-[3,5-dichloro-4-(3-bromo-4,4,4-  
trifluorobut-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxy-  
20 acrylate (compound Nr.118);

- methyl (E)-2-{2-[3,5-dichloro-4-(3-bromo-4,4,4-  
trifluorobut-2-enyloxy)phenoxy)methyl]phenyl}-2-methoxy-  
iminoacetate (compound Nr.119);

- (E)-2-{2-[3,5-dichloro-4-(3-bromo-4,4,4-trifluoro-  
25 but-2-enyloxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxy-

iminoacetamide (compound Nr.120);

- methyl (E)-2-{2-[3,5-dichloro-4-(3,4,4,4-tetrafluoro-but-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.121);

5 - methyl (E)-2-{2-[3,5-dichloro-4-(3,4,4,4-tetrafluoro-but-2-enyloxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate (compound Nr.122);

- (E)-2-{2-[3,5-dichloro-4-(3,4,4,4-tetrafluoro-but-2-enyloxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxyimino-  
10 acetamide (compound Nr.123);

- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dibromoprop-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.124);

- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dibromoprop-2-enyloxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate (com-  
15 pound Nr.125);

- (E)-2-{2-[3,5-dichloro-4-(3,3-dibromoprop-2-enyloxy)phenoxy)methyl]phenyl}-N-2-methoxyiminoacetamide (compound Nr.126);

20 - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-difluoroprop-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.127);

- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-difluoroprop-2-enyloxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate (com-  
25 pound Nr.128);

- (E)-2-{2-[3,5-dichloro-4-(3,3-difluoroprop-2-enyloxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.129);
- methyl (E)-2-{2-[5-(2,2-dichlorocyclopropyl)methoxy-  
5 2,4-dichlorophenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.130);
- methyl (E)-2-{2-[5-(2,2-dichlorocyclopropyl)methoxy-2,4-dichlorophenoxy)methyl]phenyl}-2-methoxyiminoacetate (compound Nr.131);
- 10 - (E)-2-{2-[5-(2,2-dichlorocyclopropyl)methoxy-2,4-dichlorophenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.132);
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-2,6-dichlorophenoxy)methyl]phenyl}-3-methoxyacrylate (com-  
15 pound Nr.133);
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-2,6-dichlorophenoxy)methyl]phenyl}-2-methoxyiminoacetate (compound Nr.134);
- (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-2,6-  
20 dichlorophenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.135);
- methyl (E)-2-{2-[(2,4-dichloro-5-(3,3-dichloroprop-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate (compound Nr.136);
- 25 - methyl (E)-2-{2-[(2,4-dichloro-5-(3,3-dichloroprop-2-

enyloxy)phenoxyethyl]phenyl}-2-methoxyiminoacetate (compound Nr.137);

- (E)-2-{2-[(2,4-dichloro-5-(3,3-dichloroprop-2-enyloxy)phenoxyethyl]phenyl}-N-methyl-2-methoxyimino-  
5 acetamide (compound Nr.138);

- methyl (E)-2-{2-[(2,6-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxyethyl]phenyl}-3-methoxyacrylate (compound Nr.139);

- methyl (E)-2-{2-[(2,6-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxyethyl]phenyl}-2-methoxyiminoacetate (compound Nr.140);  
10

- (E)-2-{2-[(2,6-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxyethyl]phenyl}-N-methyl-2-methoxyiminoacetamide (compound Nr.141);

15 - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy)phenoxyethyl]phenyl}-3-methoxyacrylate, m.p. 80°C,  
<sup>1</sup>H-NMR(CDCl<sub>3</sub>): δ 1.0 (9H,s), 1.8 (2H,m), 3.73 (3H,s), 3.82 (3H,s), 3.98 (2H,t), 4.93 (2H,s), 6.84 (2H,s), 7.1-7.55 (4H,m), 7.62 (1H,s); (compound Nr.142);

20 - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy)phenoxyethyl]phenyl}-2-methoxyiminoacetate, thick oil, (compound Nr. 143);

- (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy)phenoxyethyl]phenyl}-N-methyl-2-methoxyiminoacetamide,  
25 (compound Nr. 144);

- methyl (E)-2-{2-[3,5-dichloro-4-(2,4-dichlorobenzoyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate thick oil, (compound Nr. 145);

5 methyl (E)-2-{2-[3,5-dichloro-4-(4-chlorobenzoyloxy)-phenoxy)methyl]phenyl}-3-methoxyacrylate, solid with m.p. 112°C, (compound Nr. 146).

#### EXAMPLE 5

Determination of the acaricidal activity.

a) Activity against adults of *Tetranychus urticae*.

10 Samples of bean leaves are infested with adult female mites and sprayed, by means of a Potter Tower, with a hydro-acetone solution at 20% by volume of acetone, containing the product being tested at the desired concentration and Tween 20 (0.05%).

15 The mortality percentage is determined 48 hours after treatment, compared with that of adult mites, by infesting samples only treated with a hydro-acetone solution at 20% by volume of acetone (blank).

The compounds tested provided a full activity at a  
20 dose of 500 ppm.

b) Activity against eggs and larvae of *Tetranychus urticae*.

Samples of bean leaves, on which mite eggs were deposited 24 hours before, were sprayed by means of a Potter  
25 Tower with a hydro-acetone solution at 20% by volume

of acetone, containing the product being tested at the desired concentration and Tween 20 (0.05%).


The percentage of unopened eggs is determined seven days after treatment, compared with that of eggs only  
5 treated with a hydro-acetone solution at 20% by volume of acetone (blank).

72 hours after examining the eggs, the mortality percentage of the larvae deriving from the treated eggs is evaluated and compared with that of the larvae deriv-  
10 ing from the blank eggs.

The compounds tested provided a full activity at a dose of 200 ppm.

## CLAIMS

1. Compounds having general formula (I)



(I)

5 wherein:

- a group selected from  $X_1$ ,  $X_2$  and  $X_3$  represents an R group;
- $X_4$  and  $X_5$  and two of the remaining  $X_1$ ,  $X_2$ ,  $X_3$  groups represent a hydrogen atom or a halogen atom, on the con-
- 10 dition that at least two of said groups represent a halo-
- gen atom;
- R represents a  $C_1$ - $C_{12}$  alkyl or haloalkyl group; a  $C_1$ - $C_{12}$  alkoxy or alkylthio group optionally substituted by halo-
- 15 gen atoms, cyano groups,  $C_1$ - $C_6$  alkoxy groups optionally
- halogenated,  $C_2$ - $C_{10}$  alkoxyalkoxy groups optionally halo-
- genated,  $\text{NH}_2$  groups optionally substituted by  $C_1$ - $C_6$  alkyl
- groups optionally halogenated,  $C_3$ - $C_{12}$  trialkyl silyl
- groups, aryloxy or heteroaryloxy groups, in turn option-
- ally substituted by halogen atoms,  $C_1$ - $C_6$  alkyl groups op-
- 20 tionally halogenated,  $C_1$ - $C_6$  alkoxy groups optionally

halogenated, nitro groups, cyano groups; a C<sub>2</sub>-C<sub>12</sub> alkenyloxy or alkenylthio group optionally substituted by halogen atoms, cyano groups, aryl groups or heteroaryl groups, in turn optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, halo-alkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a C<sub>3</sub>-C<sub>12</sub> alkinyloxy or alkynylthio group optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkoxy or haloalkoxy groups, aryl or heteroaryl groups, in turn optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, haloalkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a linear or branched C<sub>3</sub>-C<sub>12</sub> alkoxyiminoalkylidenoxy or alkoxyiminoalkylidenthio group; a C<sub>3</sub>-C<sub>8</sub> cycloalkoxy or cycloalkylthio group optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, haloalkyl, alkoxy, haloalkoxy groups; a C<sub>4</sub>-C<sub>12</sub> cycloalkylalkoxy or cycloalkylalkylthio group optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl, haloalkyl, alkoxy, haloalkoxy groups; a heterocyclyloxy, heterocyclylthio, heterocyclyl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or heterocyclyl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio group, optionally substituted by halogen atoms, (C<sub>1</sub>-C<sub>6</sub>) alkyl, haloalkyl, alkoxy, haloalkoxy groups; an aryloxy, arylthio, hetero-aryloxy, heteroarylthio, aryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio, heteroaryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or heteroaryl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio group optionally substituted by halogen atoms, C<sub>1</sub>-C<sub>6</sub> alkyl groups optionally

halogenated, C<sub>1</sub>-C<sub>6</sub> alkoxy groups optionally halogenated, nitro groups, cyano groups;

- A, the same or different, when n is greater than or equal to 2, represents a halogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl, haloalkyl, alkoxyl, haloalkoxyl group;
- Y represents an OCH<sub>3</sub> group, an NHCH<sub>3</sub> group, an NH<sub>2</sub> group;
- Z represents a CH group or a nitrogen atom N;
- n is an integer ranging from 0 to 4.

10 2. The compounds according to claim 1, characterized in that they are an isomeric mixture in any proportion, or the isomer E or the isomer Z of the compounds having formula (I).

3. The compounds according to claim 1, characterized in  
15 that they are the isomer E of the compounds having formula (I).

4. The compounds according to claim 1, characterized in that X<sub>3</sub> represents an R group according to the above mentioned meanings, X<sub>2</sub> and X<sub>4</sub> represent a halogen atom, X<sub>1</sub>  
20 and X<sub>5</sub> represent a hydrogen atom and n is equal to 0.

5. The compounds according to claim 1, characterized in that they are selected from:


- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-3-methoxyacrylate;
- 25 - methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophen-

- oxymethyl]phenyl]-2-methoxyiminoacetate;
- (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy-methyl]phenyl]-N-methyl-2-methoxyiminoacetamide;
  - methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxy-methyl]phenyl}-3-methoxyacrylate;
  - 5 - methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxy-methyl]phenyl}-2-methoxyiminoacetate;
  - (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacet-
  - 10 amide;
  - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxy-methyl]phenyl}-3-methoxyacrylate;
  - methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxy-methyl]phenyl}-2-methoxyiminoacetate;
  - 15 - (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)-phenoxy-methyl]phenyl}-N-methyl-2-methoxyiminoacetamide;
  - methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxy-methyl]phenyl}-3-methoxy-
  - acrylate;
  - 20 - methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-tri-fluorobut-2-enyloxy)phenoxy-methyl]phenyl}-2-methoxyimi-
  - noacetate;
  - (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxy-methyl]phenyl}-N-methyl-2-methoxyimino-
  - 25 acetamide;

- methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichloro-  
phenoxy)methyl]phenyl]-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy)  
phenoxy)methyl]phenyl}-3-methoxyacrylate;
- 5 - methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy) phe-  
noxy)methyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichloro-  
phenoxy)methyl]phenyl]-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(2,4-dichloro-  
10 benzyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(4-chloro-  
benzyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate.

6. The process for the preparation of the compounds  
having general formula (I), according to any of the  
15 claims 1-5, characterized in that it includes a condensa-  
tion reaction of a compound having general formula (II)  
with a phenol having general formula (III), according to  
the reaction scheme 1:

Scheme 1



wherein ,  $X_1$ ,  $X_2$ ,  $X_3$ ,  $X_4$ ,  $X_5$ ,  $A$ ,  $Y$ ,  $Z$  and  $n$  have the meanings defined above,  $L$  represents a leaving group such as a chlorine atom, a bromine atom or a  $R_LSO_3^-$  group wherein  
 5  $R_L$  represents a  $C_1$ - $C_6$  alkyl or haloalkyl, or a phenyl optionally substituted.

7. The process according to claim 6, characterized in that the reaction is carried out in an inert organic solvent, at a temperature ranging from  $0^\circ\text{C}$  and the boiling  
 10 temperature of the reaction mixture, possibly in the presence of an inorganic or organic base.


8. The process according to claim 7, characterized in that the solvent is selected from alcohols, ethers, esters, ketones, chlorinated hydrocarbons, aromatic hydrocarbons, aliphatic hydrocarbons, aprotic dipolar solvents.  
 15

9. The process according to claim 7, characterized in that the inorganic base is selected from hydrides, hy-

droxides, carbonates of alkaline or alkaline-earth metals.

10. The process according to claim 7, characterized in that the organic base is selected from pyridine, dimethylaminopyridine, aliphatic amines, cyclic amines, alcoholates of alkaline metals.

11. Use of the compounds having general formula (I)



(I)

10 wherein:

- a group selected from  $\text{X}_1, \text{X}_2$  and  $\text{X}_3$  represents an R group;
- $\text{X}_4$  and  $\text{X}_5$  and two of the remaining  $\text{X}_1, \text{X}_2, \text{X}_3$  groups represent a hydrogen atom or a halogen atom, on the condition that at least two of said groups represent a halogen atom;
- R represents a  $\text{C}_1\text{-C}_{12}$  alkyl or haloalkyl group; a  $\text{C}_1\text{-C}_{12}$  alkoxy or alkylthio group optionally substituted by halogen atoms, cyano groups,  $\text{C}_1\text{-C}_6$  alkoxy groups optionally halogenated,  $\text{C}_2\text{-C}_{10}$  alkoxyalkoxy groups optionally halo-

generated,  $\text{NH}_2$  groups optionally substituted by  $\text{C}_1\text{-C}_6$  alkyl groups optionally halogenated,  $\text{C}_3\text{-C}_{12}$  trialkyl silyl groups, aryloxy or heteroaryloxy groups, in turn optionally substituted by halogen atoms,  $\text{C}_1\text{-C}_6$  alkyl groups optionally halogenated,  $\text{C}_1\text{-C}_6$  alkoxy groups optionally halogenated, nitro groups, cyano groups; a  $\text{C}_2\text{-C}_{12}$  alkenyloxy or alkenylthio group optionally substituted by halogen atoms, cyano groups, aryl groups or heteroaryl groups, in turn optionally substituted by halogen atoms,  $\text{C}_1\text{-C}_6$  alkyl, haloalkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a  $\text{C}_3\text{-C}_{12}$  alkinyloxy or alkynylthio group optionally substituted by halogen atoms,  $\text{C}_1\text{-C}_6$  alkoxy or haloalkoxy groups, aryl or heteroaryl groups, in turn optionally substituted by halogen atoms,  $\text{C}_1\text{-C}_6$  alkyl, haloalkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a linear or branched  $\text{C}_3\text{-C}_{12}$  alkoxyiminoalkylidenoxy or alkoxyiminoalkylidenthio group; a  $\text{C}_3\text{-C}_8$  cycloalkoxy or cycloalkylthio group optionally substituted by halogen atoms,  $\text{C}_1\text{-C}_6$  alkyl, haloalkyl, alkoxy, halo-alkoxy groups; a  $\text{C}_4\text{-C}_{12}$  cycloalkylalkoxy or cycloalkylalkylthio group optionally substituted by halogen atoms,  $\text{C}_1\text{-C}_6$  alkyl, haloalkyl, alkoxy, halo-alkoxy groups; a heterocyclyloxy, heterocyclylthio, heterocyclyl( $\text{C}_1\text{-C}_6$ )alkoxy or hetero-cyclyl( $\text{C}_1\text{-C}_6$ )alkylthio group, optionally substituted by halogen atoms, ( $\text{C}_1\text{-C}_6$ ) alkyl, haloal-

kyl, alkoxy, halo-alkoxy groups; an aryloxy, arylthio, heteroaryloxy, heteroarylthio, aryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio, heteroaryl-(C<sub>1</sub>-C<sub>6</sub>)alkoxy or heteroaryl-(C<sub>1</sub>-C<sub>6</sub>)alkylthio group optionally substituted by halogen  
5 atoms, C<sub>1</sub>-C<sub>6</sub> alkyl groups optionally halogenated, C<sub>1</sub>-C<sub>6</sub> alkoxy groups optionally halogenated, nitro groups, cyano groups;

- A, the same or different, when n is greater than or equal to 2, represents a halogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl, haloalkyl, alkoxyl, haloalkoxy group;  
10

- Y represents an OCH<sub>3</sub> group, an NHCH<sub>3</sub> group, an NH<sub>2</sub> group;

- Z represents a CH group or a nitrogen atom N;

- n is an integer ranging from 0 to 4;

15 as acaricides and/or insecticides and/or fungicides.

12. The use according to claim 11 of the isomers E of the compounds having formula (I).

13. The use according to claim 11, wherein X<sub>3</sub> represents an R group according to the above meanings, X<sub>2</sub> and X<sub>4</sub>  
20 represent a halogen atom, X<sub>1</sub> and X<sub>5</sub> represent a hydrogen atom and n is equal to 0.

14. The use according to claim 11, wherein the compounds of formula (I) are selected from:

- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophen-  
25 oxyethyl)phenyl]-3-methoxyacrylate;

- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy)methyl]phenyl]-2-methoxyiminoacetate;
- (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxy-methyl)phenyl]-N-methyl-2-methoxyiminoacetamide;
- 5 - methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxy)methyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxy)methyl]phenyl}-2-methoxyiminoacetate;
- (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-di-  
10 chlorphenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacet-  
amide;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-  
enyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-  
15 enyloxy)phenoxy)methyl]phenyl}-2-methoxyiminoacetate;
- (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)-  
phenoxy)methyl]phenyl}-N-methyl-2-methoxyiminoacetamide;
- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-  
trifluorobut-2-enyloxy)phenoxy)methyl]phenyl}-3-methoxy-  
20 acrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-tri-  
fluorobut-2-enyloxy)phenoxy)methyl]phenyl}-2-methoxyimi-  
noacetate;
- (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-tri-fluorobut-  
25 2-enyloxy)phenoxy)methyl]phenyl}-N-methyl-2-methoxyimino-

acetamide;

- methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichloro-phenoxy)methyl]phenyl]-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy) phenoxy)methyl]phenyl}-3-methoxyacrylate;
- 5 - methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy) phe-noxy)methyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichloro-phenoxy)methyl]phenyl]-3-methoxyacrylate;
- 10 - methyl (E)-2-{2-[3,5-dichloro-4-(2,4-dichloro-benzyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(4-chloro-benzyloxy)phenoxy)methyl]phenyl}-3-methoxyacrylate.

15 15. The use according to any of the claims 11-14 for the control of adults, larvae and eggs of mites and insects which are harmful in the agrarian, civil and zoo-technical field.

16. The use according to claim 15, wherein the harmful mites and/or insects are tetranychidae (*Tetranychus ur-*  
20 *ticae*, *Tetranychus telarius*, *Tetranychus cinnabarinus*, *Eotetranychus carpini*, *Panonychus ulmi*, *Panonychus citri*, etc.), eriophyidae (*Phytoptus avellanae*, *Eriophyes vitis*, *Eriophyes piri*, etc.) tarsonemidae (*Steneotarsonemus pal-*  
25 *lidus*, etc.), hemiptera (*Macrosiphum euphorbiae*, *Aphis fabae*, *Myzus persicae*, etc.), lepidoptera (*Spodoptera*

spp., *Heliothis* spp., *Chilo* spp., *Carpocapsa pomonella*, etc.), coleoptera (*Leptinotarsa decemlineata*, *Phaedon cochleariae*, etc.), diptera (*Aedes* spp., *Culex* spp., *Musca* spp., etc.).

5 17. The use according to any of the claims 11-14 for the control of phytopathogenous fungi such as: *Helminthosporium* spp., *Erysiphe* spp., *Puccinia* spp., *Plasmopara viticola*, *Pythium* spp., *Phytophthora* spp., *Rhynchosporium* spp., *Septoria* spp., *Sphaerotheca fuliginea*, *Podosphaera*  
10 *leucotricha*, *Pyricularia oryzae*, *Uncinula necator*, *Venturia* spp., *Botrytis cinerea*, *Fusarium* spp., *Alternaria* spp., *Cercospora* spp.

18. The use according to any of the claims 11-14 for the control of mites, insects and fungi which are harmful in  
15 crops of agrarian and horticultural interest, on domestic and breeding animals, in environments frequented by human beings.

19. A method for controlling mites and/or insects and/or phytopathogenous fungi in crops of agrarian and horticultural interest, and/or on domestic and breeding animals,  
20 and/or in environments frequented by human beings, by the application of the compounds having general formula (I) according to one of the claims 1-5.

20. The method according to claim 19, characterized in  
25 that the quantity of compound to be applied varies from

10 g to 5 kg per hectare.

21. Acaricidal and/or insecticidal and/or fungicidal compositions containing as active principle one or more compounds having general formula (I) according to one of the claims 1-5.

22. The compositions according to claim 21, comprising other active principles compatible with the compounds having general formula (I), such as other acaricides/insecticides, fungicides, phyto-regulators, antibiotics, herbicides, fertilizers.

23. The compositions according to claim 21, characterized in that the concentration of active principle ranges from 1 to 90%, preferably from 5 to 50%.

## INTERNATIONAL SEARCH REPORT

PCT/EP 03/03784

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C69/734 C07C259/10 A01N37/50 C07C67/343

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 145 980 A (LORENZ GISELA ET AL) 8 September 1992 (1992-09-08)  column 1, line 5 - column 3, line 35	1-3, 6-12, 15-23
Y	table 1, compounds N°1.197 to 1.202 column 31, line 43 - column 33, line 54; claims 1-3	4, 5, 13, 14
X	US 5 545 664 A (KIRSTGEN REINHARD ET AL) 13 August 1996 (1996-08-13)  column 10, line 1 - line 40 column 54, line 59 - column 55, line 15 column 66, line 35 - column 67, line 18	1-3, 6-12, 15-23
Y	table A, compounds n° A.1937-A.1958 column 84, line 33 - column 91, line 23; claims 1, 4-10	4, 5, 13, 14

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

18 August 2003

Date of mailing of the international search report

27/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Seelmann, M

## INTERNATIONAL SEARCH REPORT

PCT/EP 03/03784

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 398 692 A (SHIONOGI & CO) 22 November 1990 (1990-11-22) cited in the application formula (I-3) on page 5; scheme 4 on page 13 claims 2,14-16 -----	1-23

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1(part)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1(part)

The compounds of general formula (I) in claim 1 are defined such that for the parameter  $n$  equal or greater than 2, the radical A represents a halogen atom or a C1-C4 alkyl, haloalkyl, alkoxyl, haloalkoxyl group. The parameter  $n$  is an integer ranking from 0 to 4. For the case, wherein  $n = 1$ , no definition is provided for the radical A. There is no additional information provided in the description, which could allow to apprehend the extent of the protection sought for the radical A when  $n = 1$ .

Accordingly the search was performed only on those parts, which are supported, i.e.  $n = 0, 2, 3$  or  $4$ .

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

## INTERNATIONAL SEARCH REPORT

PCT/EP 03/03784

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5145980	A	08-09-1992	DE 3917352 A1	29-11-1990
			AU 632124 B2	17-12-1992
			AU 5589990 A	27-09-1990
			CA 2014781 A1	27-11-1990
			EP 0400417 A1	05-12-1990
			HU 54276 A2	28-02-1991
			IL 94083 A	30-03-1995
			JP 3017052 A	25-01-1991
			NZ 233815 A	29-01-1992
			ZA 9004024 A	29-01-1992
US 5545664	A	13-08-1996	AT 166341 T	15-06-1998
			AU 680718 B2	07-08-1997
			AU 1504095 A	05-10-1995
			CA 2145234 A1	26-09-1995
			CN 1121510 A	01-05-1996
			DE 59502205 D1	25-06-1998
			EP 0673923 A1	27-09-1995
			JP 7278084 A	24-10-1995
			NZ 270787 A	28-05-1996
			US 5696161 A	09-12-1997
EP 0398692	A	22-11-1990	AT 141589 T	15-09-1996
			AT 145891 T	15-12-1996
			AU 628972 B2	24-09-1992
			AU 5508890 A	22-11-1990
			BR 1100783 A3	06-06-2000
			BR 1100785 A3	21-03-2000
			BR 9002311 A	06-08-1991
			CA 2017076 A1	17-11-1990
			DE 69028136 D1	26-09-1996
			DE 69028136 T2	23-01-1997
			DE 69029334 D1	16-01-1997
			DE 69029334 T2	30-04-1997
			DK 398692 T3	16-09-1996
			DK 629609 T3	23-12-1996
			EP 0398692 A2	22-11-1990
			EP 0629609 A1	21-12-1994
			ES 2093013 T3	16-12-1996
			ES 2097583 T3	01-04-1997
			GR 3021270 T3	31-01-1997
			GR 3022142 T3	31-03-1997
			JP 2818143 B2	30-10-1998
			JP 8253449 A	01-10-1996
			JP 2759069 B2	28-05-1998
			JP 8231310 A	10-09-1996
			JP 2558001 B2	27-11-1996
			JP 3246268 A	01-11-1991
			KR 9502600 B1	23-03-1995
			US 5371223 A	06-12-1994
			US 5401877 A	28-03-1995
			US 5548078 A	20-08-1996
			US 5185342 A	09-02-1993
			US 5371222 A	06-12-1994
			ZA 9003689 A	27-02-1991

**THIS PAGE BLANK (USPTO)**